

AMPROCHEM 15

(Water Soluble Powder)

Composition:

Each 1 gm contains:

Amprolium HCl: 150 mg

Properties & Indications:

- **Amprochem 15:** is indicated for prevention and treatment of coccidiosis in broiler, layer and turkey caused by different type of Eimeria (E.tenella, E.maxima, E.acervulina, E.necatrix, E.brunetti).
- **Amprochem 15:** is also used for prevention and treatment of coccidiosis in cattle, calves, sheep and goats.

Dosage:

Poultry:

- Prevention: 4 gm / 4-8 liters of drinking water for 5 days.
- Treatment: 4 gm / 2-4 liters of drinking water for 5-7 days.

In feed: 4 kg / ton feed.

Other animals: 20-40 mg Amprolium HCl/kg b.wt for 5-7 days.

Withdrawal Period:

Don't use animal's products during treatment:

Eggs: 2 days from last treatment.

Meat/Milk: 4 days from last treatment.

Storage:

- Store in dark and dry place at temperature not exceeding 30 °C.

Package:

100gm 500gm



ADVANCED AGROCHEMICALS AND VETERINARY
PRODUCTS INDUSTRIAL CO.

Tel : 962 6 4022640

Fax: 962 6 4022340

P.O. Box: 294 Industrial Estate – Sahab / Amman -Jordan

E-mail: hussein@chemvetjo.com